

Adam J. Katz et al
U.S. Serial No.: 10/797,371
Filed: March 9, 2004
Page 2

RECEIVED
CENTRAL FAX CENTER
NOV 21 2006

In the claims:

For the Examiner's convenience, Applicants present a complete listing of the claims as submitted on November 6, 2006 with status indicator in compliance with the practice guidelines for making amendments under 37 C.F.R. §1.121(c)(1).

Additionally, please add new claims 181 and 182 as follows:

1-38. (CANCELLED)

39. (CURRENTLY AMENDED) A method of differentiating an an-clonally isolated adipose-derived stem cell into cells of any two or more of a fat cell, a bone cell, a cartilage cell, and a muscle cell comprising culturing the cell in any of adipogenic, osteogenic, chondrogenic, and myogenic morphogenic medium under conditions sufficient for the cell to differentiate.

40. (CURRENTLY AMENDED) The method of claim 39, wherein the medium is any of adipogenic, chondrogenic, cardiogenic, dermatogenic, embryonic, fetal, myogenic, osteogenic, or stromogenic media.

41. (PREVIOUSLY PRESENTED) The method of claim 39, wherein the morphogenic medium is an adipogenic medium and the cell is monitored to identify adipogenic differentiation.

42. (PREVIOUSLY PRESENTED) The method of claim 39, wherein the morphogenic medium is a chondrogenic medium and the cell is monitored to identify chondrogenic differentiation.

43. (CANCELLED)

Adam J. Katz et al
U.S. Serial No.: 10/797,371
Filed: March 9, 2004
Page 3

44. (PREVIOUSLY PRESENTED) The method of claim 39, wherein the morphogenic medium is a myogenic medium and the cell is monitored to identify myogenic differentiation.
45. (PREVIOUSLY PRESENTED) The method of claim 39, wherein the morphogenic medium is an osteogenic medium and the cell is monitored to identify osteogenic differentiation.
46. (CANCELLED)
47. (PREVIOUSLY PRESENTED) The method of claim 39, wherein the cell differentiates in vitro.
48. (PREVIOUSLY PRESENTED) The method of claim 39, wherein the cell differentiates in vivo.
- 49-56. (CANCELLED)
57. (PREVIOUSLY PRESENTED) The method of claim 39, wherein the cell is within a defined cell population.
- 58-159. (CANCELLED)
160. (CURRENTLY AMENDED) A method of inducing the differentiation of an clonally isolated adipose derived stem cell into cells of any two or more of a fat cell, a bone cell, a cartilage cell, and a muscle cell comprising culturing the cell in any of adipogenic, osteogenic, chondrogenic, and myogenic morphogenic suitable

Adam J. Katz et al
U.S. Serial No.: 10/797,371
Filed: March 9, 2004
Page 4

RECEIVED
CENTRAL FAX CENTER
NOV 21 2006

medium effective to induce differentiation under suitable differentiation conditions.

161. (PREVIOUSLY PRESENTED) The method of claim 160 wherein said medium is a conditioned medium of a specific cell type.
162. (PREVIOUSLY PRESENTED) The method of claim 160, further comprising co-culturing the cell with a cell of desired lineage.
- 163-168. (CANCELLED)
169. (NEW) The method of claim 39, wherein the muscle cell is any of skeletal muscle, cardiac muscle or smooth muscle.
170. (NEW) The method of claim 39, wherein the muscle cells express muscle specific proteins.
171. (NEW) The method of claim 170, wherein the muscle specific proteins are myoD or myosin heavy chain.
172. (NEW) The method of claim 39, wherein the cell differentiates into a precursor of a fat cell, a bone cell, a cartilage cell or a muscle cell.
173. (NEW) The method of claim 39, wherein any of adipogenic, osteogenic, chondrogenic, and myogenic morphogenic medium is supplemented with growth factors, cytokines or extracellular matrix materials.
174. (NEW) The method of claim 39, further comprising introducing the cells into a biologically compatible lattice.

Adam J. Katz et al
U.S. Serial No.: 10/797,371
Filed: March 9, 2004
Page 5

175. (NEW) The method of claim 174, wherein the lattice comprises a mesh, sponge, or hydrogel.
176. (NEW) The method of claim 174, wherein the lattice is homopolymeric or heteropolymeric.
177. (NEW) The method of claim 174, wherein the lattice comprises hormones.
178. (NEW) The method of claim 174, wherein the lattice comprises extracellular matrix materials such as fibronectin, laminin and collagen.
179. (NEW) The method of claim 57, wherein the cell is within a heterogeneous population of clonally isolated cells.
180. (NEW) The method of claim 57, wherein the cell is within a homogeneous population of clonally isolated cells.
181. (NEW) The method of claim 39, wherein differentiation is determined by morphological changes, staining with cell specific stains, biochemical markers indicative of the desired phenotype, expression of cell specific molecules, sorting by size and granularity, or by antibody binding to cell specific molecules.
182. (NEW) The method of claim 161, wherein the conditioned medium is medium to which the clonally isolated adipose derived stem cells are exposed and cultured.